


COZIFLEX

Undenatured Type II Collagen 40 mg



Composition :

Each delayed release (DR) capsule contains Undenatured Type II Collagen 40mg.

Description :

Undenatured Type II Collagen derived from chicken sternum cartilage using a patented, low-temperature, non-enzymatic manufacturing process that preserves the natural triple helix molecular configuration and its biological activity. Undenatured Type II Collagen has unique mechanism of action. Undenatured Type II Collagen molecules contain active building sites called epitopes which trigger the process of oral tolerization. Through oral tolerization process, active epitopes of Undenatured Type II Collagen interact with Peyer's patches (Lymphoid tissue) of small intestine to release collagen-specific regulatory T-cells. Collagen-specific regulatory T-cells then migrate to joint areas and prevent release of T-cells. Preventing the release of T-cells prevent the secretion of collagenase enzymes by macrophages responsible for breakdown of collagen in joint cartilage. It also slows the production of inflammatory cytokines resulting in reduction of inflammation, halting the erosion of joint cartilage and promote cartilage rebuilding.

COZIFLEX Supports joint function, mobility and flexibility and also improves knee extension by up to 10% for better range of motion. It is also relieves joint pain due to strenuous exercise.

Indications :

COZIFLEX capsule is indicated for :

- ① Rheumatoid Arthritis
- ① Osteoarthritis
- ① Joint pain & inflammation
- ① Joint discomfort & stiffness (knee extension)
- ① Strenuous exercise & sports
- ① Difficulties in flexibility & mobility
- ① Impaired joint function
- ① Inflammation associated with injury

Dosage and administration :

Orally take 1 capsule daily with meal. Or, as directed by the registered physician.

Contraindication :

It is contraindicated in those patients with known hypersensitivity to chicken or egg.

Precautions :

There is no data available.

Side Effects :

Undenatured Type II Collagen is generally well tolerated in recommended dose.

Use in Pregnancy and Lactation :

There are no adequate & well controlled studies in pregnant and lactating women.

Drug Interactions :

There is no data available.

Over dosage :

Overdose may cause constipation and headache.

Storage :

Store below 30°C in a cool and dry place. Keep out of the reach of children. Keep away from direct sunlight.

Packing :

Each box contains 30 delayed release (DR) capsule in blister pack.

Manufactured by



DRUG INTERNATIONAL LTD.

Herbal Division

Tongi, Gazipur, Bangladesh

05/2025/Nil